



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

May 6, 2014

N. Bhushan Mandava, Ph.D., Agent
Repar Corporation
c/o Mandava Associates, LLC
1050 Connecticut Ave., NW, Suite 1000
Washington, DC 20036

Subject: Label Amendments (change ingredients statement, create sub-labels for
consumer and professional uses)
Repar Weed and Feed
EPA Reg. No. 69361-2
Resubmission Dated March 14, 2014

Dear Dr. Mandava:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

A stamped copy of the label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products released for shipment after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", written over a horizontal line.

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs



Master Label
REPAR WEED & FEED

Net Wt. 17.5 Lbs. (7.93 kg)
Treats Up To 20,000 Sq. Ft.

This label contains two separate sub-labels for, **Residential Label (Sub-Label A)** and **Commercial Label (Sub-Label B)** product use.

- **For use on warm and cool season turf types as listed.**
- **Controls dandelions, clover, dollarweed, chickweed and 200 common lawn weeds (as listed).**
- **For a thick green weed free lawn.**

ACTIVE INGREDIENTS:

Chlorflurenol –Methyl	0.69%
Dicamba	0.13%

OTHER INGREDIENTS:..... 99.18%

Total.....100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Manufactured for:
Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914

A C C E P T E D
05/06/2014
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 69361-2

EPA Reg. No. 69361-2

EPA Est. No.-

Fertilizer and Carrier information will be supplied by registrants/sub-registrants to States and other regulatory authorities who have jurisdiction over these ingredients.



SUB-LABEL A: RESIDENTIAL LABEL REPAR WEED & FEED

Net Wt. 17.5 Lbs. (7.93 kg)
Treats Up To 20,000 Sq. Ft.

- For use on warm and cool season turf types as listed.
- Controls dandelions, clover, dollarweed, chickweed and 200 common lawn weeds (as listed).
- For a thick green weed free lawn.

ACTIVE INGREDIENTS:

Chlorflurenol -Methyl	0.69%
Dicamba	0.13%
OTHER INGREDIENTS.....	99.18%
Total.....	100.00%

EPA Reg. No. 69361-2

EPA Est. No. -

Fertilizer and Carrier information will be supplied by registrants/sub-registrants to States and other regulatory authorities who have jurisdiction over these ingredients.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) call **CHEMTREC** at 1-800-424-9300.

For PRODUCT USE Information call (202) 223 – 1424.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt and long pants, socks, shoes and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

SEE REVERSE SIDE FOR ADDITIONAL INFORMATION

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide of the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

WEED CONTROL

Repar Weed and Feed controls a wide range of annual, biennial, perennial and woody weeds including:

WEED LIST

Annuals

Amaranth, Spiny (Spiny Pigweed)	Goosefoot, Nettleleaf	Pennycress, Field (Fanweed,	Senna, Coffee
Aster, Slender	Henbil	Frenchweed,	Sesbania, Hemp
Bedstraw	Jimsonweed	Stinkweed)	Shephardspurse
Beggarweed, Florida	Knotweed	Pepperweed, Virginia (Peppergrass)	Sicklepod
Broomweed, Common	Kochia	Pigweed, Prostrate	Sida, Prickly (Teaweed)
Buckwheat, Wild	Kochia (triazine resistant)	Pigweed, Redroot (Carelessweed)	Smartweed, Green
Buffalobur	Ladysthumb	Pigweed, Rough (Carelessweed)	Smartweed, Pennsylvaniz
Burclover, California	Lambsquarters, Common	Pigweed, Rough (triazine resistant)	Sneezeweed, Bitter
Burcucumber	Lambsquarters, (triazine resistant)	Pigweed, Tumble	Sowthistle, Annual
Buttercup, Roughseed	Lettuce, Prickly	Poorjoe	Sowthistle, Spiny
Carpetweed	Mallow, Common	Puncturevine	Spanishneedles
Catchfly, Nighflowering	Mallow, Venice	Purslane, Common	Spikeweed, Common
Chamomile, Corn	Mareslail (Horseweed)	Pusley, Florida	Spurge
Chickweed, Common	Mayweed	Radish, Wild	Spurr, Corn
Clovers, Annual	Medic, Black	Ragweed, Common	Starbur, Bristly
Cockle, Corn	Morningglory, Ivyleaf	Ragweed, Giant (Buffaloweed)	Stumpweed, Rough
Cocklebur, Common	Morningglory, Tall	Ragweed, Lance-Leaf	Sunflower, Common (Wild)
Croton, Tropic	Mustard, Tansy	Rubberweed, Bitter (Bitterweed)	Sunflower, Volunteer
Croton, Woolly	Mustard, Wild		Thistle, Russian
Daisy, English	Mustard (Yellowtaps)		Velvetleaf
Eveningprimrose,	Nightshade, Black		Waterhemp
Cutleaf			Waterprimrose, Winged
Fleabane, Annual			Wormwood, Annual

Biennials

Burdock, Common	Geranium	Plantain, Bracted	Thistle, Bull
Carrot, Wild	Gromwell	Ragwort, Tansy	Thistle, Milk
(Queen Anne's Lace)	Knapweed, Diffuse	Starthistle, Yellow	Thistle, Musk
Cockle, White	Knapweed, Spotted	Sweetclover	Thistle, Plumelless
Eveningprimrose, Common	Mallow, Dwarf	Teasel	

Perennials

Alfalfa*	Clover, Hop*	Milkweed, Climbing	Sowthistle
Artichoke, Jerusalem	Dandelion, Common*	Milkweed, Common	Sowthistle, Perennial
Aster, Spiny	Dock, Broadleaf*	Milkweed, Honeyvine	Spurge, Leafy
Aster, Whitehead	(Bitterdock)	Milkweed, Western	Sundrop, Halfshrub
Bedstraw, Smooth	Dock, Curly*	Whorled	(Eveningprimrose)
Bindweed, Field	Dogbane, Hemp	Nettle, Stinging	Thistle., Canada
Bindweed, Hedge	Dogfennel*	Nightshade, Silverleaf	Toadflax Dalmation
Bindweed, Texas	(Cypressweed)	(White Horsenettle)	Tropical Soda Apple
Bursage*	Fern, Bracken	Onion, Wild	Trumpetcreeper
(Bur Ragweed,	Garlic, Wild	Pennywort (Dollarweed)	(Buckvine)
Lakeweed,	Goldenron, Canada	Plantain, Broadleaf*	Vetch
Povertyweed)	Goldenrod, Missouri	Plantain, Buckhorn	Violet, Wild
Bursage, Woolyleaf	Goldenweed, Common	Pokeweed	Waterhemlock
(Lakeweed)	Hawkweed	Ragweed, Western	Waterprimrose
Buttercup, tall	Henbane, Black	Sericia Lespedeza	Creeping
Campion, Bladder	Horsenettle, Carolina	Redvine	Woodsorrel, Creeping*
Chickweed, Field	Ironweed	Smartweed, Swamp	(Common Yellow)
Chickweed, Mouseear	Ivy, Ground	Snakeweed, Broom	Wormwood, Common
(Canada)	Knapweed, Black	Sorel, Red*	Wormwood, Louisiana
Chicory	Knapweed, Russian	(Sheep Sorel)	Yankeeweed*
			Yarrow, Common

*Noted perennials may be controlled using Repair Weed and Feed at rates lower than those directed for other listed perennial weeds.

Woody

Ailanthus	Ivy, Poison	Kudzu	Sagebrush, Fringe
Sassfras	Gallberry	Serviceberry	Wild Grape
Oak, Poison	Sumac	Brazilian Pepper	Tarbrush
Honeysuckle	Hornbeam	Witchhazel	Huckleberry
Huisache			

For Use on Established Turfs, including sensitive species such as Centipede and St. Augustine, except Desirable Clovers, Bent, Dichondra and Carpet Grass.

EFFECTIVE APPLICATION OF REPAIR WEED AND FEED FOR MOST WEED LEAVES SHOULD BE DAMP FROM RAIN, DEW OR SPRINKLING TO ALLOW THE PRODUCT TO STICK TO THE LEAF. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL.

WHAT TO LOOK FOR:

Repair Weed and Feed, properly applied, first reduces or eliminates the formation of most flower heads and the seeds they produce and then attacks and removes the mature weed itself. Look for an absence or stunting of seed heads in flowering weeds such as clover or dandelions. Some weeds may also show twisting stems and dwarfed appearance with curling of the leaves.

Repair Weed and Feed is designed with an excellent turf nutrient diet to allow grass plants to gradually fill in the areas around dying weeds. This allows for a more natural recovery of the turf and minimizes reinfestations of bare areas by wind-borne weed seeds. Applied as directed, the rich green color will not be affected.

UNDER NORMAL GROWING CONDITIONS, WEED KILL SHOULD BE COMPLETE WITHIN 21 TO 35 DAYS

SPREADER SETTINGS

Spreader	Rate: 3.5#/M	Pattern Setting	Width
Earthway Broadcast	11	-	8
Lely	4 1/2	III - B	42
Lesco Broadcast	D 1/2	0.0	11
Prizelawn Broadcast	4 1/4	A	7
Scotts Broadcast	1	5	10
Spyker	3	-	9
Vicon	29	-	4
Cyclone Broadcast	3 3/4	-	5

Important: These are approximate settings. Age and condition of your spreader can cause wide variation. Be sure to calibrate your spreader with each application.

WHEN TO APPLY:

Apply any time weeds are actively growing and temperature is above 60°F. Do not apply when turf is under stress from heat, excessive moisture or drought. Control is quicker and more complete if applied in Spring prior to weed flower formation. Control is slower if applied when weather is cool. On newly seeded areas, wait until grass has been cut two or three times before using. Delay planting new grass for at least three weeks as the active ingredients may temporarily inhibit seed germination.

HOW TO APPLY:

EFFECTIVE APPLICATION OF REPAIR WEED AND FEED REQUIRES THE WEED LEAVES TO BE DAMP FROM RAIN, DEW OR SPRINKLING PRIOR TO APPLICATION. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL. Apply evenly over the entire lawn at the rate of 0.875 lbs./1,000 sq. ft. Do not make more than 2 applications per year. Rain or heavy sprinkling with 24 hours of application will temporarily delay weed control. Mowing within 24 hours may reduce effectiveness. Wait at least 45 days before making a second application.

Use the suggested spreader settings as a guide being sure to check your spreader at the beginning of the application. Adjust your spreader if necessary. Be sure to shut off spreader while turning or stopping to prevent accidental over application and burning of grass. SPECIAL NOTE: Repair Weed And Feed is designed to be absorbed by plant leaves. Accidental application of this product to the foliage of ornamentals such as roses may result in the loss of flowers. For this reason, extra care should be taken when using a broadcast or rotary type spreader.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep container closed and away from food, feedstuffs and domestic water supplies.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

IMPORTANT:

READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. To the extent consistent with applicable law, manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS: YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION.

Manufactured for:
Repair Corporation, P.O. Box 4321
Silver Spring, Maryland 20914



SUB-LABEL B: COMMERCIAL LABEL REPAR WEED & FEED

Net Wt. 17.5 Lbs. (7.93 kg)
Treats Up To 20,000 Sq. Ft.

- For use on warm and cool season turf types as listed.
- Controls dandelions, clover, dollarweed, chickweed and 200 common lawn weeds (as listed).
- For a thick green weed free lawn.

ACTIVE INGREDIENTS:

Chlorflurenol –Methyl	0.69%
Dicamba	0.13%
OTHER INGREDIENTS:	99.18%
Total.....	100.00%

EPA Reg. No. 69361-2

EPA Est. No.-

Fertilizer and Carrier information will be supplied by registrants/sub-registrants to States and other regulatory authorities who have jurisdiction over these ingredients.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) call **CHEMTREC** at 1-800-424-9300.

For PRODUCT USE Information call (202) 223 – 1424.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves made of any waterproof material.

SEE REVERSE SIDE FOR ADDITIONAL INFORMATION

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any irrigation system.

Do not apply this product in a way that will contact any person, or animals, either directly or through drift.

Only protected handlers may be in the area during application.

Do not allow people or animals to enter the treated area until dusts have settled.

WEED CONTROL

Repar Weed and Feed controls a wide range of annual, biennial, perennial and woody weeds including:

WEED LIST

Annuals

Amaranth, Spiny (Spiny Pigweed)	Goosefoot, Nettleleaf	Pennycress, Field (Fanweed,	Senna, Coffee
Aster, Slender	Jimsonweed	Frenchweed,	Sesbania, Hemp
Bedstraw	Knotweed	Stinkweed)	Shephardspurse
Beggarweed, Florida	Kochia	Pepperweed, Virginia (Peppergrass)	Sicklepod
Broomweed, Common	Kochia (triazine resistant)	Pigweed, Prostrate	Sida, Prickly (Teaweed)
Buckwheat, Wild	Ladysthumb	Pigweed, Redroot (Carelessweed)	Smartweed, Green
Buffalobur	Lambsquarters, Common	Pigweed, Rough (Carelessweed)	Smartweed, Pennsylvaniz
Burclover, California	Lambsquarters, (triazine resistant)	Pigweed, Rough (triazine resistant)	Sneezeweed, Bitter
Burcucumber	Lettuce, Prickly	Pigweed, Tumble	Sowthistle, Annual
Buttercup, Roughseed	Mallow, Common	Poorjoe	Sowthistle, Spiny
Carpetweed	Mallow, Venice	Puncturevine	Spanishneedles
Catchfly, Nighflowering	Mareslail (Horseweed)	Purslane, Common	Spikeweed, Common
Chamomile, Corn	Mayweed	Pusley, Florida	Spurge
Chickweed, Common	Medic, Black	Radish, Wild	Spurr, Corn
Clovers, Annual	Morningglory, Ivyleaf	Ragweed, Common	Starbur, Bristly
Cockle, Corn	Morningglory, Tall	Ragweed, Giant (Buffaloweed)	Stumpweed, Rough
Cockle, Cow	Mustard, Tansy	Ragweed, Lance-Leaf	Sunflower, Common (Wild)
Cocklebur, Common	Mustard, Wild	Rubberweed, Bitter (Bitterweed)	Sunflower, Volunteer
Croton, Tropic	Mustard (Yellowtaps)		Thistle, Russian
Croton, Woolly	Nightshade, Black		Velvetleaf
Daisy, English			Waterhemp
Eveningprimrose,			Waterprimrose, Winged
Cutleaf			Wormwood, Annual
Fleabane, Annual			

Biennials

Burdock, Common	Geranium	Plantain, Bracted	Thistle, Bull
Carrot, Wild (Queen Anne's Lace)	Gromwell	Ragwort, Tansy	Thistle, Milk
Cockle, White	Knapweed, Diffuse	Starthistle, Yellow	Thistle, Musk
Eveningprimrose, Common	Knapweed, Spotted	Sweetclover	Thistle, Plumeless
	Mallow, Dwarf	Teasel	

Perennials

Alfalfa*	Clover, Hop*	Milkweed, Climbing	Sowthistle
Artichoke, Jerusalem	Dandelion, Common*	Milkweed, Common	Sowthistle, Perennial
Aster, Spiny	Dock, Broadleaf*	Milkweed, Honeyvine	Spurge, Leafy
Aster, Whiteheath	(Bitterdock)	Milkweed, Western	Sundrop, Halfshrub (Eveningprimrose)
Bedstraw, Smooth	Dock, Curly*	Whorled	Thistle., Canada
Bindweed, Field	Dogbane, Hemp	Nettle, Stinging	Toadflax Dalmation
Bindweed, Hedge	Dogfennel*	Nightshade, Silverleaf	Tropical Soda Apple
Bindweed, Texas	(Cypressweed)	(White Horsesettle)	Trumpetcreeper
Bursage*	Fern, Bracken	Onion, Wild	(Buckvine)
(Bur Ragweed, Lakeweed, Povertyweed)	Garlic, Wild	Pennywort (Dollarweed)	Vetch
Bursage, Woolyleaf (Lakeweed)	Goldenron, Canada	Plantain, Broadleaf*	Violet, Wild
Buttercup, tall	Goldenrod, Missouri	Plantain, Buckhorn	Waterhemlock
Campion, Bladder	Goldenweed, Common	Pokeweed	Waterprimrose
Chickweed, Field	Hawkweed	Ragweed, Western	Creeping
Chickweed, Mouseear (Canada)	Henbane, Black	Sericia Lespedeza	Woodsorrel, Creeping* (Common Yellow)
Chicory	Horsenettle, Carolina	Redvine	Wormwood, Common
	Ironweed	Smartweed, Swamp	Wormwood, Louisiana
	Ivy, Ground	Snakeweed, Broom	Yankeeweed*
	Knapweed, Black	Sorrel, Red*	Yarrow, Common
	Knapweed, Russian	(Sheep Sorrel)	

*Noted perennials may be controlled using Repair Weed and Feed at rates lower than those directed for other listed perennial weeds.

Woody

Ailanthus	Ivy, Poison	Kudzu	Sagebrush, Fringe
Sassfras	Gallberry	Serviceberry	Wild Grape
Oak, Poison	Sumac	Brazilian Pepper	Tarbrush
Honeysuckle	Hornbeam	Witchhazel	Huckleberry
Huisache			

For Use on Established Turfs, including sensitive species such as Centipede and St. Augustine, except Desirable Clovers, Bent, Dichondra and Carpet Grass.

EFFECTIVE APPLICATION OF REPAIR WEED AND FEED FOR MOST WEED LEAVES SHOULD BE DAMP FROM RAIN, DEW OR SPRINKLING TO ALLOW THE PRODUCT TO STICK TO THE LEAF. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL.

WHAT TO LOOK FOR:

Repair Weed and Feed, properly applied, first reduces or eliminates the formation of most flower heads and the seeds they produce and then attacks and removes the mature weed itself. Look for an absence or stunting of seed heads in flowering weeds such as clover or dandelions. Some weeds may also show twisting stems and dwarfed appearance with curling of the leaves.

Repair Weed and Feed is designed with an excellent turf nutrient diet to allow grass plants to gradually fill in the areas around dying weeds. This allows for a more natural recovery of the turf and minimizes reinfestations of bare areas by wind-borne weed seeds. Applied as directed, the rich green color will not be affected.

UNDER NORMAL GROWING CONDITIONS, WEED KILL SHOULD BE COMPLETE WITHIN 21 TO 35 DAYS

SPREADER SETTINGS

Spreader	Rate: 3.5#/M	Pattern Setting	Width
Earthway Broadcast	11	-	8
Lely	4 1/2	III - B	42
Lesco Broadcast	D 1/2	0.0	11
Prizelawn Broadcast	4 1/4	A	7
Scotts Broadcast	1	5	10
Spyker	3	-	9
Vicon	29	-	4
Cyclone Broadcast	3 3/4	-	5

Important: These are approximate settings. Age and condition of your spreader can cause wide variation. Be sure to calibrate your spreader with each application.

WHEN TO APPLY:

Apply any time weeds are actively growing and temperature is above 60°F. Do not apply when turf is under stress from heat, excessive moisture or drought. Control is quicker and more complete if applied in Spring prior to weed flower formation. Control is slower if applied when weather is cool. On newly seeded areas, wait until grass has been cut two or three times before using. Delay planting new grass for at least three weeks as the active ingredients may temporarily inhibit seed germination.

HOW TO APPLY:

EFFECTIVE APPLICATION OF REPAIR WEED AND FEED REQUIRES THE WEED LEAVES TO BE DAMP FROM RAIN, DEW OR SPRINKLING PRIOR TO APPLICATION. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL. Apply evenly over the entire lawn at the rate of 0.875 lbs./1,000 sq. ft. Do not make more than one application per year. Not for use on sod farms. Rain or heavy sprinkling with 24 hours of application will temporarily delay weed control. Mowing within 24 hours may reduce effectiveness.

Use the suggested spreader settings as a guide being sure to check your spreader at the beginning of the application. Adjust your spreader if necessary. Be sure to shut off spreader while turning or stopping to prevent accidental over application and burning of grass. SPECIAL NOTE: Repair Weed And Feed is designed to be absorbed by plant leaves. Accidental application of this product to the foliage of ornamentals such as roses may result in the loss of flowers. For this reason, extra care should be taken when using a broadcast or rotary type spreader.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep container closed and away from food, feedstuffs and domestic water supplies.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. To the extent consistent with applicable law, manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS: THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS: YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION.

Manufactured for:
Repair Corporation, P.O. Box 4321
Silver Spring, Maryland 20914